


PICODIV

Monitoring the diversity of photosynthetic picoplankton in marine waters

EU contract EVK3-CT-1999-00021



Objectives: Picoplankton (defined operationally as cells that pass through a 3 micron filter) dominate the photosynthetic biomass in many marine ecosystems, not only in the very oligotrophic regions of the world oceans, such as the Eastern Mediterranean Sea, but also in mesotrophic areas. However, picophytoplankton are clearly not exclusively restricted to pelagic environments. In many coastal regions, they are present throughout the year and constitute a 'background' population, onto which episodic phenomena such as the spring bloom develops. Fewer than 50 species of picophytoplankton have been described. A clear proof of our poor knowledge of picophytoplankton diversity is revealed by the discovery of three novel algal classes in the last ten years described from picophytoplanktonic taxa. Because so little is known about the taxonomy and systematics of picophytoplankton we have very little data to estimate the levels of its biodiversity under natural conditions and how picophytoplankton are affected by environmental variability linked to either anthropogenic influence or to larger scale phenomena such as those linked to climate change or global warming.

The major objective of the PICODIV was to develop, test and validate probing methods based on molecular biology techniques that allow for routine and extensive assessment of picophytoplankton diversity (species composition and relative contribution of taxa to total community) in the marine environment. Our strategy to meet this objective was encapsulated in the following four steps: (1) Obtain SSU rDNA sequences for as many as possible picophytoplankton taxa from both cultures and natural samples. Novel taxa will be assessed using a combination of methods including in particular pigment analysis and electron microscopy. (2) Using this sequence database, develop hierarchical probes recognizing each taxonomic group having picophytoplanktonic representatives (3) Develop fast and efficient techniques to quantify the fraction of the pico-phytoplankton recognized by the probes in natural samples. (4) Test and validate these probes on time series of picophytoplankton biodiversity in three coastal ecosystems.

Scientific achievements.

Cultures of picoplankton.

Obtaining cultures of novel picoplanktonic species is a key step to assess picoplankton diversity. More than 250 cultures have been established from three coastal sites (Roscoff, Helgoland, Blanes) and from open ocean cruises. These cultures have been characterized by microscopy, pigment analysis and rDNA sequencing. We performed a detailed analysis of the phenotype (pigments, morphology) and genotype (18S rDNA phylogeny) of key groups of the picoplankton based on representative strains isolated during the project. This work led to the reorganization of the taxonomy of important groups such as the cyanobacteria and Prasinophyceae as well as to the description of novel

species (e.g. *Florenciella parvula*) or the correct phylogenetic assignment of previously described species (e.g. *Telonema*).

Clone libraries of marine picoplankton.

Construction of SSU rDNA clone libraries representative of the *in situ* picophytoplankton diversity, and subsequent sequencing of novel gene sequences, is critical for understanding and describing the total diversity present in contrasting marine aquatic environments. This is particularly so for identification of novel picophytoplankton groups for which cultured counterparts do not exist. The sequence data obtained provide the basis for the design of new phylogenetic probes.

In the course of the project we have obtained over 1,300 rDNA sequences from 46 clone libraries constructed both from coastal and open ocean samples. We analyzed the sequence data in order to both determine the spatial and seasonal variation of taxonomic groups at each coastal site and perform a detailed phylogenetic reconstruction for some of the groups, especially those for which no culture representatives are available (stramenopiles, alveolates).

Molecular probes for marine picoplankton.

We have designed and tested the specificity of probes against some key taxonomic groups from the picoplankton. We then used probes to monitor the abundance in coastal waters of different groups among picoplankton (*Synechococcus*, Prasinophyceae, Prymnesiophyceae...) with detection techniques developed during the project, in particular fluorescent *in situ* hybridization coupled with the tyramide signal amplification (FISH-TSA) and DNA micro-arrays.

Socio-economic relevance and policy implications: The collection of cultures that have been isolated during the project should be a major resource both from a fundamental point of view because these strains are likely to represent key organisms in coastal waters and from the applied point of view because these strains could be screened in the future for interesting compounds. From the methodological point of view, the DGGE approach, that we have developed and applied, could prove very useful to assess and monitor the diversity of phytoplankton in oceanic waters. Finally the set of probes we have designed and tested as well the probing techniques (FISH-TSA, DNA micro-arrays) that have been developed should be very useful for routine monitoring of some key groups, such as the *Synechococcus* cyanobacteria and the Prasinophyceae, e.g. in the context of long term surveys to detect ecosystem response to global change or to pollution.

Conclusions: The PICODIV project has led to major advances in the knowledge of the diversity of picoplankton in coastal waters, revealing the ubiquity of novel groups that had been recently discovered in open ocean waters (stramenopiles, alveolates). Moreover, we have been able to establish the precise seasonal dynamics of key photosynthetic groups such as cyanobacteria and Prasinophyceae, using molecular tools and to relate these estimates to those made from well established methods such as pigments analysis.

Selected references from PICODIV

- Biegala, I., Not, F., Vault, D. & Simon, N. 2003. Quantitative assessment of picoeucaryotes in the natural environment using taxon specific oligonucleotide probes in association with TSA-FISH (Tyramide Signal Amplification - Fluorescent In Situ Hybridization) and flow cytometry. **Applied and Environmental Microbiology** in press.
- Díez, B., Pedrós-Alió, C., Marsh, T. L. & Massana, R. 2001. Application of denaturing gradient gel electrophoresis (DGGE) to study the diversity of marine picoeukaryotic assemblages and comparison of DGGE with other molecular techniques. **Applied and Environmental Microbiology** 67:2942-2951.
- Díez, B., Pedrós-Alió, C. & Massana, R. 2001. Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing. **Applied and Environmental Microbiology** 67:2932-2941.
- Fuller, N. J., Marie, D., Partensky, F., Vault, D., Post, A. F. & Scanlan, D. J. 2003. Clade-specific 16S ribosomal DNA oligonucleotides reveal the predominance of a single marine *Synechococcus* clade throughout a stratified water column in the Red Sea. **Applied and Environmental Microbiology** 69:2430-2443.
- Massana, R., Guillou, L., Díez, B. & Pedros Alio, C. 2002. Unveiling the organisms behind novel eukaryotic ribosomal DNA sequences from the ocean. **Applied and Environmental Microbiology** 68:4554-4558.
- Moon-van der Staay, S. Y., De Wachter, R. & Vault, D. 2001. Oceanic 18S rDNA sequences from picoplankton reveal unsuspected eukaryotic diversity. **Nature** 409:607-610.
- Vault, D., Romari, K. & Not, F. 2002. Are autotrophs less diverse than heterotrophs in marine picoplankton? **Trends in Microbiology** 10:266-267.
- West, N. J., Schonhuber, W. A., Fuller, N. J., Amann, R. I., Rippka, R., Post, A. F. & Scanlan, D. J. 2001. Closely related *Prochlorococcus* genotypes show remarkably different depth distributions in two oceanic regions as revealed by *in situ* hybridization using 16S rRNA-targeted oligonucleotides. **Microbiology** 147:1731-1744.